

Experimental prospects for indirect BSM searches in $e^+e^- \rightarrow q\bar{q}$ ($q=b,c$) processes at Higgs Factories.

SW&ANA Meeting 13/03/2024

J. P. Márquez, A. Irles, A. Saibel, H. Yamamoto – AITANA Group at IFIC (CSIC/UV) - Valencia

F. Richard, R. Poesch – IJCLab - Orsay

N. Yamatsu – National Taiwan University



Gen=T



CSIC

CONSEJO SUPERIOR DE INVESTIGACIONES CIENTÍFICAS

IFIC

INSTITUT DE FÍSICA
CORPUSCULAR



VNIVERSITAT
DE VALÈNCIA



Financiado por
la Unión Europea
NextGenerationEU



GENERALITAT
VALENCIANA
Conselleria de Educació,
Universitats y Empleo



Plan de Recuperación,
Transformación y Resiliencia



Irène Joliot-Curie
Laboratoire de Physique
des 2 Infinis



National
Taiwan
University



AITANA

Introduction & motivation




Motivation: BSM Z' resonances

► Many **BSM scenarios** (i.e. Randal Sundrum, compositeness, Gauge Higgs unification models...) predict heavy resonances coupling to the (t,b) doublet and also lighter fermions (i.e. c/s quarks)

- Only coupling to (t,b) doublet
 - → Peskin, Yoon arxiv:1811.07877
 - → Djouadi et al arxiv:hep-ph/0610173
- Coupling also to lighter fermions [Funatsu, Hatanaka, Hosotani, Orikasa, Yamatsu]

(arxiv:1705.05282) (arxiv:2309.01132) (arxiv:2301.07833)




Probing such scenarios require at least per mil level for experimental precision

tt/bb/cc/ss/... **Can we do it?**

Gauge-Higgs Unification Models (GHU)

- ▶ Randall-Sundrum metric (5D).
- ▶ The symmetry breaking pattern is different than in the SM and features the so-called Hosotani's mechanism.
- ▶ Only one parameter, Hosotani's angle θ_H , determines the projection of the 5D fields, fixing all physical effects:
 - KK resonances of the Z/ γ with $m_{kk} \sim 10\text{-}25$ TeV.
 - Modifications and new EW couplings/helicity amplitudes.
 - Already visible effects at 250GeV.

As **Benchmark**, we will use the [Funatsu, Hatanaka, Hosotani, Orikasa, Yamatsu] models.



▶ A models: ([arxiv:1705.05282](https://arxiv.org/abs/1705.05282))

$$A_1 : \theta_H = 0.0917, m_{KK} = 8.81 \text{ TeV} \rightarrow m_{Z^1} = 7.19 \text{ TeV};$$

$$A_2 : \theta_H = 0.0737, m_{KK} = 10.3 \text{ TeV} \rightarrow m_{Z^1} = 8.52 \text{ TeV},$$

▶ B models: ([arxiv:2309.01132](https://arxiv.org/abs/2309.01132)) ([arxiv:2301.07833](https://arxiv.org/abs/2301.07833))

$$B_1^\pm : \theta_H = 0.10, m_{KK} = 13 \text{ TeV} \rightarrow m_{Z^1} = 10.2 \text{ TeV};$$

$$B_2^\pm : \theta_H = 0.07, m_{KK} = 19 \text{ TeV} \rightarrow m_{Z^1} = 14.9 \text{ TeV};$$


$$B_3^\pm : \theta_H = 0.05, m_{KK} = 25 \text{ TeV} \rightarrow m_{Z^1} = 19.6 \text{ TeV};$$



GHU vs SM (250 GeV)


$\sqrt{s_{e^+e^-}} = 250 \text{ GeV}$

b-quark



$\sqrt{s_{e^+e^-}} = 250 \text{ GeV}$

c-quark




$$A_{FB} = \frac{N_F - N_B}{N_F + N_B}$$

GHU vs SM (500 GeV)


$\sqrt{s_{e^+e^-}} = 500 \text{ GeV}$

b-quark



$\sqrt{s_{e^+e^-}} = 500 \text{ GeV}$

c-quark




$$A_{FB} = \frac{N_F - N_B}{N_F + N_B}$$

GHU vs SM (1 TeV)


$\sqrt{s_{e^+e^-}} = 1000 \text{ GeV}$

b-quark



$\sqrt{s_{e^+e^-}} = 1000 \text{ GeV}$

c-quark



$$A_{FB} = \frac{N_F - N_B}{N_F + N_B}$$

Experimental study with full simulation





- ▶ ILD note and previous works <https://inspirehep.net/literature/2669897>
 - ILC250, b and c studies. (A. Irles, F. Richard, R. Poesch).
- ▶ Work presented in LCWS (J.P. Márquez):
 - Proceeding <https://inspirehep.net/literature/2682331>
 - Talk: https://indico.slac.stanford.edu/event/7467/contributions/5977/attachments/2862/8042/LCWS2023_JPMH.pdf
 - ILC250+ILC500 comparing scenarios with different PID (no PID, dEdx, dNdx).
 - Experimental cut-based analysis using “traditional” BDT algorithms for flavour tagging.
- ▶ Work presented in EPS-HEP (J.P. Márquez).
 - Proceeding: <https://inspirehep.net/literature/2714494>
 - Talk: <https://indico.desy.de/event/34916/contributions/147224/>
 - First theory prospects.
- ▶ Update presented at ECFA in Paestum (A. Irles).
 - Final theory prospects.
- ▶ Presented at the last ILD meeting at CERN (J.P. Márquez).

Jet flavour tagging & charge measurement





► Double tagging & charge measurement methods. (described in previous [ILD note](#))

- To maximally reduce the usage of MC tools (control of fragmentation, QCD correlations... uncertainties)




Result and fit

- ▶ **At least 4 observables for AFB at ILC per energy point**
 - 2 quarks (b and c).
 - 2 polarizations (e_{LpR} , e_{RpL}).
- ▶ **Per mil level statistical uncertainties reachable for the nominal ILC program**
- **Smaller exp syst. Uncertainties**
- Fragmentation, angular correlations, preselection efficiency...



Result ILC250 & ILC500



Discrimination power between GHU & SM




- ▶ Assumption: A measurement of one specific model is conducted.
 - ▶ The uncertainties are considered normally distributed:
 - Significance in σ .
 - P-value: Gaussian at d_σ .
- $$d_\sigma = \frac{\|AFB_{\text{test}} - AFB_{\text{ref}}\|}{\Delta_{AFB_{\text{ref}}}}$$
- ▶ Combination of multiple measurements is done with a *multivariate gaussian*.
 - Assuming no correlations for AFB.





GHU vs SM: discrimination power plots

ILD
work in progress



GHU vs SM: GHU energy scale

B_3^+		m_{KK}
B_3^-		25 TeV
B_2^+		25 TeV
B_2^-		19 TeV
B_1^+		13 TeV
B_1^-		13 TeV
A_2		m_{KK}
A_1		10.3 TeV
		8.8 TeV

GHU vs SM: Beam scenarios

Hypothetical case
ILC250* no pol
 $\int L = 2000 \text{ fb}^{-1}$

**Full ILD simulation
assuming
no beam pol.**

H20 nominal program

ILC250
($P_{e-}=0.8, P_{e+}=0.3$)
 $\int L = 2000 \text{ fb}^{-1}$


ILC500
($P_{e-}=0.8, P_{e+}=0.3$)
 $\int L = 4000 \text{ fb}^{-1}$

**Full ILD simulation
assuming beam pol.**


H20 nominal program

ILC1000
($P_{e-}=0.8, P_{e+}=0.2$)
 $\int L = 8000 \text{ fb}^{-1}$

*Not full simulation studies
but extrapolations from ILC500*




GHU vs SM: c. m. e.



GHU vs SM: Precision on Z-couplings

GHU vs SM discrimination power (σ -level)


ILD
work in progress



GHU vs SM: Precision on Z-couplings


ILD
work in progress

GHU vs SM discrimination power (σ -level)



GHU vs SM: Beam(s) polarization

work in progress





GHU vs SM: Positron beam polarization

work in progress




GHU vs SM discrimination power (σ -level)



GHU vs SM: Positron beam polarization

work in progress





GHU vs SM: Particle ID dependence



work in progress

GHU vs SM discrimination power (σ -level)




GHU vs SM: Particle ID dependence

work in progress




GHU vs SM discrimination power (σ -level)







GHU vs SM: Particle ID dependence

work in progress

GHU between model discrimination



Conclusion/ summary



- ▶ Comprehensive study done at ILC250/ILC500 with ILD simulations:
 - Backgrounds, beam features, polarization, realistic reconstruction tools.
 - Uncertainties dominated by statistics, above the Z-pole.
 - Room for improvement (modern algorithms for flavour tagging, event selection, etc).
- ▶ AFB studies for c and b-quark above the Z-pole.
 - Flavour tagging and jet charge determination with kaon ID are key.
- ▶ ILC offers unique capabilities to explore these signatures and discriminate GHU vs SM:
 - High energy reach.
 - Electron and positron beam polarization → enhancing the sensitivity but also allowing for measurements with different BSM sensitivity (for control of systematics)
- (ILD) PID capabilities (kaon)



- ▶ A paper has been prepared
 - Ready for submission for EPJ-C.
- ▶ ILD editorial board:
 - Reviewed by Mikael Berggren and Daniel Jeans.
 - Circulated on the ILD mailing list.
- ▶ Everything set for journal review!

Eur. Phys. J. C manuscript No.
(will be inserted by the editor)

Probing Gauge-Higgs Unification models at the ILC with di-quark forward-backward asymmetry at center-of-mass energies above the Z mass. ^{*}

A. Irles^{1,a}, J.P. Márquez¹, R. Pöschl², F. Richard², A. Saibel¹,
H. Yamamoto^{b,1}, N. Yamatsu³

¹IFIC, Universitat de València and CSIC. C./ Catedrático José Beltrán 2, E-46100 Paterna, Spain

²Université Paris-Saclay, CNRS/IN2P3, IJCLab, 91405 Orsay, France

³Department of Physics, National Taiwan University, Taipei, Taiwan 10617, R.O.C.

Received: date / Accepted: date

Abstract The International Linear Collider (ILC) will allow the precise study of $e^+e^- \rightarrow q\bar{q}$ interactions at different center-of-mass energies from the Z -pole to 1 TeV. In this paper we discuss the experimental prospects for measuring differential observables in $e^+e^- \rightarrow b\bar{b}$ and $e^+e^- \rightarrow c\bar{c}$ at the ILC baseline energies, 250 and 500 GeV. The studies are based on full detector simulation samples and reconstruction of the International Large Detector (ILD) concept. Two gauge-Higgs unification models predicting new high-mass resonances beyond the Standard Model are discussed. These models predict sizable deviations of the forward-backward observables at the ILC running above the Z mass and with longitudinally polarized electron and positron beams. The ability of the ILC to probe these models via high-precision forward-backward asymmetry measurements is discussed. Alternative scenarios with other energy points or different beam polarisation schemes are also discussed, extrapolating the estimated uncertainties from the two baseline scenarios.

Keywords First keyword · Second keyword · More

1 Introduction

The Standard Model (SM) is a successful theory, well-established experimentally and theoretically. With the discovery of the Higgs boson [1, 2], the structure of the SM seems to be confirmed. However, some inconsistencies in the SM still need to be answered. For instance,

the striking mass hierarchy in the fermion sector. Moreover, while the dynamic of the SM gauge bosons, the photon, W and Z bosons, and gluons are governed by the gauge principle, the dynamic of the Higgs boson is different and unique in the SM. The SM does not predict the values of the Higgs couplings of quarks and leptons, nor the Higgs self-couplings. Large quantum corrections have to be canceled by fine-tuning the parameters to calculate the Higgs boson mass matching the measured value. One possible solution to this issue, achieving stabilization of the Higgs mass against quantum corrections, appears when the Higgs boson is associated with the zero mode of a dimension-five component of extensions of the SM gauge group. These models are referred to as gauge-Higgs unification (GHU) models.

The two most precise determinations of $\sin^2 \theta_{eff}$ by the LEP and SLC differ in 3.7σ , and none of them agrees with the SM prediction [3, 3]. In particular, the LEP value was extracted from the forward-backward asymmetry measurement for b -quarks with LEP1 data, and it is nearly three standard deviations away from the predicted value in the SM. Clarifying the A_{FB}^b value as well as exploring the possibility of BSM physics motivate the study of quark pair production in high energy e^-e^+ collisions at future colliders not only at the Z -mass energy but also at higher energies. In the SM, these interactions are produced and mediated by a photon, a Z -boson, and the interference between them. Some BSM theories predict deviation of such couplings or even new sizable contributions to these processes from new mediators (such as heavy Z' resonances). These deviations would be accessible experimentally by performing high precision measurements of $e^-e^+ \rightarrow q\bar{q}$ observables at different center-of-mass energies (\sqrt{s}).

^{*}This work was carried out in the framework of the ILD concept group.

^aCorresponding author: adrian.irles@ific.uv.es

^bOn leave from Tohoku University, Sendai, Japan



Thanks for your attention!

back-up



► <https://arxiv.org/pdf/2203.07622.pdf>

Quantity	Value	current $\delta[10^{-4}]$	Z pole		ILC250	
			$\delta_{stat}[10^{-4}]$	$\delta_{sys}[10^{-4}]$	$\delta_{stat}[10^{-4}]$	$\delta_{sys}[10^{-4}]$
boson properties						
m_W	80.379	1.5	-	-	0.08	0.3
m_Z	91.1876	0.23		0.022		-
Γ_Z	2.4952	9.4	0.5	-	6	-
$\Gamma_Z(had)$	1.7444	11.5		4.	-	-
Z-e couplings						
$1/R_e$	0.0482	24.	2.	5	5.5	10
A_e	0.1513	139.	1.5	1.2	12.	9.
g_L^e	-0.632	16.	1.0	3.2	2.8	7.6
g_R^e	0.551	18.	1.0	3.2	2.9	7.6
Z- ℓ couplings						
$1/R_\mu$	0.0482	16.	2.	2.	5.5	10
$1/R_\tau$	0.0482	22.	2.	2.	5.7	10
A_μ	0.1515	991.	2.	5	54.	3.
A_τ	0.1515	271.	2.	5.	57.	3
g_L^μ	-0.632	66.	1.0	2.3	4.5	7.6
g_R^μ	0.551	89.	1.0	2.3	5.5	7.6
g_L^τ	-0.632	22.	1.0	2.8	4.7	7.6
g_R^τ	0.551	27.	1.0	3.2	5.8	7.6
Z-b couplings						
R_b	0.2163	31.	0.4	7.	3.5	10
A_b	0.935	214.	1.	5.	5.7	3
g_L^b	-0.999	54.	0.32	4.2	2.2	7.6
g_R^b	0.184	1540	7.2	36.	41.	23.
Z-c couplings						
R_c	0.1721	174.	2.	30	5.8	50
A_c	0.668	404.	3.	5	21.	3
g_L^c	0.816	119.	1.2	15.	5.1	26.
g_R^c	-0.367	416.	3.1	17.	21.	26.

GHU vs SM: Beam scenarios

Hypothetical case
ILC250* no pol
 $\int L = 2000\text{fb}^{-1}$
OSP|SSP [%] = 45 | 5
Full ILD simulation
assuming
no beam pol


H20 nominal program

ILC250
($P_{e-}=0.8, P_{e+}=0.3$)
 $\int L = 2000\text{fb}^{-1}$
OSP|SSP [%] = 45 | 5
ILC500
($P_{e-}=0.8, P_{e+}=0.3$)
 $\int L = 4000\text{fb}^{-1}$
OSP|SSP [%] = 40 | 10
Full ILD simulation
assuming beam pol.


H20 nominal program

ILC1000
($P_{e-}=0.8, P_{e+}=0.2$)
 $\int L = 8000\text{fb}^{-1}$
OSP|SSP [%] = 40 | 10

Not full simulation studies
but extrapolations from ILC500



- ▶ **Topology: 2 back-to-back jets (pencil-like topology)**
- ▶ **Preselection aiming for high background rejection and high efficiency.**
- ▶ Main bkg $ee \rightarrow Z\gamma$ (radiative return through ISR)
 - $\sim x10$ larger than signal
 - **$\sim 90\%$ of such ISR photons are lost in the beam pipe** \rightarrow events filtered by energy & angular mom. conservation arguments
 - The **remaining $\sim 10\%$ are filtered by identifying photons** in the detector (efficiency of $>90\%$)
 - PFA detector!!
- ▶ Other backgrounds from diboson production decaying hadronically are removed with extra topological cuts.



Double-Tag method

► Compare samples with 1 tag vs 2 tags (after preselection)

$$f_{1b} = \overline{\varepsilon_c} \overline{R_b} + \overline{\tilde{\varepsilon}_c} \overline{R_c} + \overline{\tilde{\varepsilon}_{uds}} (1 - \overline{R_b} - \overline{R_c})$$
$$f_{2b} = \overline{\varepsilon_b^2} (1 + \overline{\rho}) \overline{R_b} + \overline{\varepsilon_c^2} \overline{R_c} + \overline{\varepsilon_{uds}^2} (1 - \overline{R_b} - \overline{R_c})$$

Measured observables





PHYSICS!
Indirect observables

Inputs (MC or independent measurements)

**Similar set of equations
for the c-quark
solved simultaneously**

Double flavour tagging – control of systematics

- ▶ **Flavour tagging efficiency will be measured (double tagging)**
 - Not estimated with MC
 - **Per mil level reachable** because the contamination from lighter quarks is minimal and the tight IP constraint
- ▶ **Fully differential analysis !!**
- ▶ **R_b and R_c measured at the same time**
 - than the tagging efficiencies
 - No assumption needed in Ruds
- ▶ **Per mil level stat. Uncertainty**
- ▶ **Comparable/lower exp syst. uncertainty**
 - Dominated by flavour tagging and followed by angular correlations



ILD-PHYS-PUB-2023-001


▶ We start from a very pure & background-free **double tagged** sample



▶ We are required to **measure the jet charge**

- Using K-ID and/or full Vtx charge measurement
- K-ID is better suited for the C-quark (Vtx is better suited for b-quark)

▶ We use the **double charge** measurements

- To control / reduce the systematic uncertainties





ILD-PHYS-PUB-2023-001

Double charge method

► Double Tag + Double Charge


- Both jets need to have a charge measurement compatible with the 2 quarks back to back scenario

► Double mistakes are unlikely but still not negligible and lead to “sign flip” → migrations

BSM or simple migrations?

Red shows the distribution without sign correction.


Gray is the parton level distribution



ILD-PHYS-PUB-2023-001

Migration correction


- ▶ Migrations look as “new physics” → we need to correct them
 - Using data: double charge measurements with same and opposite charges (see back-up slides)
 - We measure the probability to reconstruct correctly the charge (P_b) and use it for correction
 - DATA DRIVEN METHOD → non sensitive to fragmentation modelling.



$$e^+e^- \rightarrow b\bar{b}$$
$$P_{e^+e^-} = (-0.8, +0.3) / Lum =$$

blue shows the distribution after sign correction.




Gray is the parton level distribution



▶ Pchg limited by vertex reconstruction efficiency, Particle ID efficiency and B0 oscillations (b-quark case).

ILD-PHYS-PUB-2023-001





Migration correction – c quark case



Minimal migration effects
(and corrections!)



Total uncertainties



Statistical uncertainties dominate over systematic uncertainties