

# Corrections to the MSSM Higgs Boson Mass of $\mathcal{O}(\alpha_t \alpha_s^2)$

Philipp Kant

in collaboration with  
R. V. Harlander, L. Mihaila, M. Steinhauser

Phys.Rev.Lett.100:191602,2008

Karlsruhe University

ILC-ECFA Workshop 2008

why consider Higgs sector of MSSM?

- ▶ quite predictive  
two new free parameters at tree level

$$M_A, \tan \beta$$

- ▶ important for possible discovery of MSSM  
light supersymmetric Higgs might be first trace of  
supersymmetry
- ▶ experimental accuracy  
 $\delta M_h \approx 100 - 200 \text{ MeV}$  at LHC,  $\delta M_h \approx 50 \text{ MeV}$  at ILC

## Higgs Potential

$$V = m_1^2 |H_1|^2 + m_2^2 |H_2|^2 + m_{12}^2 \left( \epsilon_{ab} H_1^a H_2^b + \epsilon_{ab} H_1^{a*} H_2^{b*} \right) + \frac{1}{8} (g_1^2 + g_2^2) \left( |H_1|^2 - |H_2|^2 \right)^2 + \frac{1}{2} g_2^2 |H_1^* H_2|^2$$

spontaneous symmetry breaking:  $H_1, H_2$  acquire vacuum expectation values  $\Rightarrow$  gauge bosons and fermions acquire masses.



difference to the SM : quartic terms fixed by gauge couplings

$$\Rightarrow M_h \text{ can be predicted, tree level: } M_h \leq M_Z$$

# Radiative Corrections to $M_h$

- ▶ 1-loop corrections from top and stop loops  $\propto M_t^4$

[Ellis,Ridolfi,Zwirner 1991; Haber,Hempfling 1991; Okada,Yamaguchi,Yanagida 1991]





significant shift of up to 35GeV

# Radiative Corrections to $M_h$

- ▶ 1-loop corrections from top and stop loops  $\propto M_t^4$

[Ellis,Ridolfi,Zwirner 1991; Haber,Hempfling 1991; Okada,Yamaguchi,Yanagida 1991]



significant shift of up to 35GeV


- ▶ from other sectors of the MSSM

[Chankowski,Pokorski,Rosiek 1994; Dabelstein 1995; Bagger,Matchev,Pierce,Zhang 1997]



small against the corrections from tops and stops

# Relevance of $M_t^4$ Approximation



# 2loop Corrections to $M_h$

- ▶ corrections in effective potential approximation ( $p^2 = 0$ )



[ Brignole, Carena, Casas, Dedes, Degrassi, Espinosa, Haber, Hempfling, Heinemeyer, Hoang, Hollik, Martin, Quirós, Riotto, Rzehak, Slavich, Wagner, Weiglein, Zhang, Zwirner, ... '94-today]

# 2loop Corrections to $M_h$

- corrections in effective potential approximation ( $p^2 = 0$ )

[ Brignole, Carena, Casas, Dedes, Degrassi, Espinosa, Haber, Hempfling, Heinemeyer, Hoang, Hollik, Martin, Quirós, Riotto, Rzehak, Slavich, Wagner, Weiglein, Zhang, Zwirner, ... '94-today]

contributions from  $t$  and  $\tilde{t}$  loops becomes even more important!





# 2loop Corrections to $M_h$

- ▶ corrections in effective potential approximation ( $p^2 = 0$ )

[ Brignole, Carena, Casas, Dedes, Degrassi, Espinosa, Haber, Hempfling, Heinemeyer, Hoang, Hollik, Martin, Quirós, Riotto, Rzehak, Slavich, Wagner, Weiglein, Zhang, Zwirner, ... '94-today]

- ▶ momentum dependence  $\approx 160\text{MeV}$

[Martin '05]

# 2loop Corrections to $M_h$

- ▶ corrections in effective potential approximation ( $p^2 = 0$ )  
[ Brignole, Carena, Casas, Dedes, Degrassi, Espinosa, Haber, Hempfling, Heinemeyer, Hoang, Hollik, Martin, Quirós, Riotto, Rzehak, Slavich, Wagner, Weiglein, Zhang, Zwirner, ... '94-today]
- ▶ momentum dependence  $\approx 160\text{MeV}$  [Martin '05]
- ▶ 3loop LL and NLL through Renormalisation Group [Martin '07]

remaining uncertainty: 3 – 5GeV  
against 100 – 200MeV at LHC, 50MeV at ILC

# Aim: Genuine Three-Loop Calculation

---

simplifying assumptions

- ▶ effective potential approximation
- ▶ restrict to  $t, \tilde{t}$  sector  
 $t, \tilde{t}, g, \tilde{g}, q$  and  $\tilde{q}$  as internal particles

# Aim: Genuine Three-Loop Calculation

simplifying assumptions

- ▶ effective potential approximation
- ▶ restrict to  $t, \tilde{t}$  sector  
 $t, \tilde{t}, g, \tilde{g}, q$  and  $\tilde{q}$  as internal particles
- ▶ assume mass hierarchy

$$M_q = 0, \quad M_t \ll M_{\tilde{t}_1} = M_{\tilde{t}_2} = M_{\tilde{g}} \ll M_{\tilde{q}}$$

asymptotic expansion in small ratios of masses

# Aim: Genuine Three-Loop Calculation

simplifying assumptions

- ▶ effective potential approximation
- ▶ restrict to  $t, \tilde{t}$  sector  
 $t, \tilde{t}, g, \tilde{g}, q$  and  $\tilde{q}$  as internal particles
- ▶ assume mass hierarchy

$$M_q = 0, \quad M_t \ll M_{\tilde{t}_1} = M_{\tilde{t}_2} = M_{\tilde{g}} \ll M_{\tilde{q}}$$

asymptotic expansion in small ratios of masses

- ▶ neglect  $A_t$  contributions  
corrections to off-diagonal elements of massmatrix vanish

# Computational Setup

---

- ▶ lots and lots of diagrams (about 16,000) ...
- ▶ ... require automatisaton: QGRAF, Q2E, EXP, MINCER, MATAD, FORM

[Nogueira; Harlander, Seidensticker; Larin, Tkachov; Steinhauser; Vermaseren]



# Computational Setup

- ▶ lots and lots of diagrams (about 16,000) ...
- ▶ ... require automatisation: QGRAF, Q2E, EXP, MINCER, MATAD, FORM

[Nogueira; Harlander, Seidensticker; Larin, Tkachov; Steinhauser; Vermaseren]

- ▶ regularisation through dimensional reduction [Siegel '79]  
 $\epsilon$ -scalars implemented,  $M_\epsilon = 0$  on-shell
- ▶ on-shell renormalisation  
recalculation of all relevant renormalisation constants

- ▶ degenerate  $\tilde{t}$  masses, no mixing between  $\tilde{t}_1$  and  $\tilde{t}_2$   
mixing angle receives radiative corrections





- ▶ degenerate  $\tilde{t}$  masses, no mixing between  $\tilde{t}_1$  and  $\tilde{t}_2$   
 mixing angle receives radiative corrections




cannot force degenerate bare  $\tilde{t}$  masses  
 $\Rightarrow$  expand integrals in  $m_{\tilde{t}_1}^2 - m_{\tilde{t}_2}^2$

- ▶ degenerate  $\tilde{t}$  masses, no mixing between  $\tilde{t}_1$  and  $\tilde{t}_2$   
 mixing angle receives radiative corrections



cannot force degenerate bare  $\tilde{t}$  masses  
 $\Rightarrow$  expand integrals in  $m_{\tilde{t}_1}^2 - m_{\tilde{t}_2}^2$

- ▶ finite infrared “divergences”




either use small  $\epsilon$ -scalar mass, or small external momentum

$$\begin{aligned}
 \Delta M_h = & -\frac{3G_F M_t^4}{\sqrt{2}\pi^2} \left\{ -L_{tS} + \frac{\alpha_s}{\pi} [4L_{tS} - 2L_{tS}^2] + \left(\frac{\alpha_s}{\pi}\right)^2 \left[ -\frac{1091}{324} - \frac{1}{27}\pi^2 - \frac{1}{9}\zeta_3 \right. \right. \\
 & + \left( \frac{1591}{108} + 3L_{\mu t} - \frac{1}{3}\pi^2 + \frac{4}{9}\pi^2 \ln 2 - \frac{55}{18}L_{t\bar{q}} - \frac{5}{6}L_{t\bar{q}}^2 \right) L_{tS} \\
 & + \left( -\frac{19}{18} - \frac{3}{2}L_{\mu t} + \frac{5}{3}L_{t\bar{q}} \right) L_{tS}^2 - \frac{53}{18}L_{tS}^3 \\
 & + \left( -\frac{475}{108} + \frac{5}{9}\pi^2 \right) L_{t\bar{q}} + \frac{25}{36}L_{t\bar{q}}^2 + \frac{5}{18}L_{t\bar{q}}^3 \\
 & \left. \left. + \mathcal{O}\left(\frac{M_S^2}{M_{\bar{q}}^2}\right) \right] \right\},
 \end{aligned}$$


$$L_{tS} = \ln \frac{M_t^2}{M_{SUSY}^2}, \quad L_{\mu t} = \ln \frac{\mu^2}{M_t^2}, \quad L_{t\bar{q}} = \ln \frac{Mt^2}{M_{\bar{q}}^2}$$

- ▶ agreement with literature
  - ▶ exact 2-loop [FeynHiggs; Degrassi, Slavich, Zwirner '01]
  - ▶ 3-loop LL and NLL [Martin '07]
- ▶ every part of the calculation has been done twice
- ▶ calculation performed in general covariant gauge  
gauge parameter independent result
- ▶ calculated also the case of unbroken SUSY  
loop corrections to  $M_h$  vanish


# Numerical Impact




# Reduced Scale Dependency



# Numerical Impact




# Numerical Impact







# Numerical Impact




# Numerical Impact




# Numerical Impact




# Full Result vs LL Approximation



# Full Result vs LL Approximation



# Full Result vs LL Approximation



- ▶ three-loop calculation of  $M_h$  in the MSSM, effect of about 500MeV
  - ▶ relevant for LHC and ILC
- ▶ renormalisation scale dependence drastically improved
- ▶ next step: allow for different mass hierarchies